

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY)

19 May 2003

2. REPORT TYPE

View Graphs

3. DATES COVERED (From - To)

4. TITLE AND SUBTITLE

New Materials Design

5a. CONTRACT NUMBER

5b. GRANT NUMBER

5c. PROGRAM ELEMENT NUMBER

6. AUTHOR(S)

Jerry Boatz, Mark Gordon, Greg Voth, Sharon Hammes-Schiffer, Ruth Pachter

5d. PROJECT NUMBER

2303

5e. TASK NUMBER

M2C8

5f. WORK UNIT NUMBER

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)

AFRL/PRSP

10 E. Saturn Blvd.

Edwards AFB CA 93524-7680

8. PERFORMING ORGANIZATION REPORT NUMBER

AFRL-PR-ED-VG-2003-138

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)

AFRL/PRS

5 Pollux Drive

Edwards AFB CA 93524-7048

10. SPONSOR/MONITOR'S ACRONYM(S)

11. SPONSOR/MONITOR'S NUMBER(S)

AFRL-PR-ED-VG-2003-138

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT

20030617 140

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:

17. LIMITATION OF ABSTRACT

18. NUMBER OF PAGES

19a. NAME OF RESPONSIBLE PERSON  
Sheila Benner

a. REPORT

b. ABSTRACT

c. THIS PAGE

Unclassified

Unclassified

Unclassified

A

19b. TELEPHONE NUMBER (include area code)  
(661) 275-5693

Standard Form 298 (Rev. 8-98)  
Prescribed by ANSI Std. Z39.18

FILE

MEMORANDUM FOR PRS (Contractor Publication)

19 May 2003

FROM: PROI (STINFO)

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-VG-2003-138

5364 Boatz, Jerry (PRSP), "New Materials Design"

DoD High Performance Computing Users Group Conference  
(Bellevue, WA, 9-13 Jun 2003) (Deadline = 9 Jun 2003)

(Statement A)

# New Materials Design

DoD UGC, 9-13 June 2003

Bellevue, WA




Jerry Boatz

Senior Research Chemist

Propulsion Directorate

Air Force Research Laboratory



# NEW MATERIALS DESIGN



## THE TEAM....

Prof. Mark S. Gordon

Prof. Gregory Voth




Prof. Sharon Hammes-Schiffer

PENNSYLVANIA




Dr. Ruth Pachter, AFRL/MLPJ



Dr. Jerry Boatz, AFRL/PRSP





# OUTLINE



## 1. Project Overview


- High energy density materials
- Polyhedral oligomeric silsesquioxanes (POSS)
- Non-linear optical materials

## 2. Theoretical Methods and benchmarks

- Ab initio electronic structure theory
- Nuclear-electronic orbital approach


## 3. Results

## 4. Summary




# PROJECT OVERVIEW - HEDM


## High Energy Density Matter -- next generation rocket propellants



High-nitrogen/polynitrogen compounds




Energetic Ionic Liquids



Specific Impulse

$$I_{sp} \propto \sqrt{\Delta H / m}$$



# PROJECT OVERVIEW - HEDM



## Technical issues being addressed using CCM

### 1. High-nitrogen/polynitrogen compounds


**Objective:** identify, characterize, and synthesize stable compounds with high heats of formation, high densities

- geometries, energy content, stabilities, reaction pathways

### 2. Energetic ionic liquids


**Objective:** identify, characterize, and synthesize thermally stable energetic ILS with low melting points, high densities and low viscosities.

- geometries, electronic structures, stabilities, energy content, interaction potentials



# PROJECT OVERVIEW - POSS

## Polyhedral oligomeric silsesquioxanes -- next generation plastics



### As Additives



Heat/abrasion resistant paints and coatings  
 Mechanical property/viscosity/thermal modifiers  
 Crosslinking agents  
 Fire retardants

### As Plastics


Medical materials  
 Space resistant resins  
 Packaging/coatings  
 Electronic materials  
 Optical Plastics

### As Preceramics

Ablative materials (nozzles, insulations etc.)  
 Claddings/electronics coatings  
 Precursors to glassy or ceramic matrices







# PROJECT OVERVIEW - POSS



## Technical issues being addressed using CCM

### 1. Mechanisms of formation

**Objective:** rational design and synthesis of POSS

- role of solvents, acid/base catalysis, substituent effects on mechanism of formation

### 2. Potential applications as molecular “sieves”


**Objective:** determine if POSS cages can be used to separate small molecules

- determine barriers to encapsulation of  $N_2$  and  $O_2$

### 3. Ti-POSS as new catalysts

**Objective:** determine if Ti-POSS and Ti-siloxane compounds are effective catalysts


- determine barriers to polymerization of ethylene, oxidation by  $HO_2H$ .




# PROJECT OVERVIEW - NLO



## Non-linear optical materials for laser-hardened applications






# PROJECT OVERVIEW - NLO




## Technical issues being addressed using CCM

### 1. Mechanism of reverse saturable absorption (RSA)



Five-level model for nonlinear absorption



# THEORETICAL METHODS

## 1. Ab initio electronic structure theory


- General Atomic and Molecular Electronic Structure System (GAMESS) -- a CHSSI code
- Nuclear-electronic orbital approach (NEO) for including nuclear quantum effects (important, e.g., in proton transfer reactions)

Various computational techniques are employed to solve the molecular electronic Schrödinger equation from quantum mechanics:

$$\left[ -\frac{1}{2} \sum_i \nabla_i^2 - \sum_i \sum_{\alpha} \frac{Z_{\alpha}}{r_{i\alpha}} + \sum_i \sum_{j>i} \frac{1}{r_{ij}} \right] \Psi_{el} = E_{el} \Psi_{el}$$

Categories of approximate solutions:

- a) “Self-consistent field” (SCF): reasonably good geometries
- b) “Electron correlation”: post-SCF correction, required for reliable energetics (e.g., barriers).



# THEORETICAL METHODS


## 1. Ab initio electronic structure theory (cont.)

- Most electronic structure codes use Born-Oppenheimer (i.e., “clamped nuclei”) approximation -- NOE method treats specified nuclei at QM level.

### Nuclear-Electronic Hamiltonian

$$\begin{aligned}
 H_{\text{tot}}(\mathbf{r}_e, \mathbf{r}_q; \mathbf{r}_c) = & -\sum_i \frac{N_e}{2} \nabla_i^2 - \sum_i \sum_A \frac{N_e N_c Z_A}{r_{iA}} + \sum_i \sum_{j>i} \frac{N_e N_e}{r_{ij}} \\
 & - \sum_I \frac{N_p}{2M_I} \nabla_I^2 + \sum_I \sum_A \frac{N_p N_c Z_A Z_I}{r_{IA}} + \sum_I \sum_{J>I} \frac{N_p N_p Z_I Z_J}{r_{IJ}} \\
 & - \sum_i \sum_I \frac{N_e N_p Z_I}{r_{iI}} + \sum_A \sum_{B>A} \frac{N_c N_c Z_A Z_B}{r_{AB}}
 \end{aligned}$$

- $N_e$ : number of electrons (coordinates  $\mathbf{r}_e$ )  
 $N_p$ : number of quantum nuclei (coordinates  $\mathbf{r}_p$ )  
 $N_c$ : number of classical nuclei (coordinates  $\mathbf{r}_c$ )



# THEORETICAL METHODS



## Ab initio electronic structure theory


### Current Status of parallel GAMESS

	<u>RHF</u>	<u>ROHF</u>	<u>UHF</u>	<u>GVB</u>	<u>MCSCF</u>
Energy	cdp	cdp	cdp	cdp	cdp
Analytic Gradient	cdp	cdp	cdp	cdp	cdp
Numeric Hessian	cdp	cdp	cdp	cdp	cdp
Analytic Hessian	cdp	cdp	-	cdp	-
MP2 energy	cdp	cdp	cdp	-	c p
MP2 gradient	cdp	-	cd	-	-
CI energy	cdp	cdp	-	cdp	cdp
CI gradient	cd	-	-	-	-
DFT energy	cdp	cdp	cdp	-	-
DFT gradient	cdp	cdp	cdp	-	-

c = conventional disk storage of AO integrals

d = direct evaluation of AO integrals

p = runs in parallel





# THEORETICAL METHODS



## Ab initio electronic structure theory

### MP2 Gradient Scalability Test






# The Search for New Polynitrogens




- All polynitrogens are unstable with respect to  $N_2$  molecules
- Their activation energy for  $N_2$  elimination is largely determined by the weakest bond in the compound
- Their metastability is enhanced if suitable resonance structures exist:



- The double-bond character of the  $\text{N}-\text{N}$  bonds in the azide anion explains its exceptional stability
- How can this stabilization effect be used to our advantage in preparing new compounds?







# Pentazolate ( $N_5^-$ )?



- Substituted pentazoles  $R-N_5$  have been known for decades ( $R=aryl$ )
- Cyclic  $N_5^-$  is aromatic
- Conversion of the diazonium salt,  $RN_2^+$ , to the substituted pentazole ring  $R-N_5$  by the reaction with azide ion,  $N_3^-$ , has been demonstrated many years ago by Ugi and Huisgen.
- $N_5^-$  has been recently detected in the gas phase for the first time, using collisional fragmentation (electrospray ion mass spectroscopy).
- Can a chemical route to  $N_5^-$  be found? (e.g., can a suitable R group be found for the reaction  $R-N_2^+ + N_3^- \rightarrow R-N_5 \rightarrow R^+ + N_5^-$ ?)





# New Polynitrogen Anions



## Pentazole anion ( $N_5^-$ )

- Theoretical calculations show that this anion has a 28 kcal/mole activation energy barrier for decomposition and its decomposition to  $N_3^-$  and  $N_2$  is only 11 kcal/mol exothermic
- Aryl substituted pentazoles can be isolated as stable compounds only if stored at low temperatures. In methanol, these compounds rapidly decompose at room temperature to form aryl azides and  $N_2$  gas






# Synthetic Challenge – How do we make These New Anions??




## Synthesis of Substituted Pentazoles


### Sources for the Pentazole Anion ( $N_5^-$ )




Trityl/Silyl Diazonium Salts



Aryl Diazonium Salts





# *Theoretical Challenge - Can we design and predict viable precursors to $N_5^-$ ?*



Find a substituent R so that

1.  $R-N_2^+$  is stable wrt  $R^+ + N_2$ .
2. The R-N bond in  $R-N_5$  is weak, thereby suitable for the reaction  $R-N_5 + M^+X^- \rightarrow M^+N_5^- + R-X$

## Approach


Use quantum chemical calculations to predict the stability of  $R-N_2^+$  diazonium salts and the length/strength of the  $R-N_5$  bond.

MBPT(2)/6-31G(d) level of theory




M = C, Si

L = F, Cl,  $CH_3$ ,  $CF_3$ ,  $NO_2$ , phenyl, etc.




## Identifying Precursors for New Polynitrogens


*This ion has been suggested  
as a useful precursor to new  
polynitrogen molecules...*



*... but calculations predict it to be  
unstable.*



Computational requirements: ~50,000 CPU-hours, 1.2 GW on IBM SP/P3 at ASC




# RESULTS - HEDM




## High-nitrogen/polynitrogen compounds

Potential dicyanoimidazolate  $[R_3Si-N_2]^+$  precursors to precursor to displacement of  $N_5^-$  formation of suitable  $R_3Si-N_5$  substituted pentazoles





$R_3$	$R_3Si-N_2^+$ distance(Å)	$D_e(Si-N_2^+)$ (kcal/mol)	$R_3Si-N_5$ distance (Å)
3Me	2.151	13.8	1.855
2Me,F	2.144	15.7	1.834
Me,2F	2.151	21.2	1.855
3F	1.973	33.4	1.783

Computational requirements: ~50,000 CPU-hours, 1 GW on IBM SP/P3 at ASC






# RESULTS - POSS

Nuclear quantum effects in water-catalyzed condensation reactions



Level	Reactive Complex	Transition State	Products	CPU Time
RHF/6-31G*	-20.4	16.5	-7.3	1.0
RHF/6-31G*/ZPE	-15.6	21.2	-7.0	
NEO/HF/4	-23.0	7.2	-5.9	1.1
NEO/HF/8	-24.8	7.7	-6.7	1.8





## POSS as catalysts?




Some Ti compounds are known to be effective catalysts

Ziegler-Natta, olefin oxidation

Recent experiments reveal Ti-modified silicates catalyze olefin oxidation by peroxides

Can Ti-substituted siloxanes and/or POSS catalyze olefin oxidation by  $\text{HOOH}$ , or polymerization?







# POSS computational methods




- § B3LYP/6-31G(d) geometry optimizations
- § Hessians to characterize stationary points
- § Energies using MP2/TZVP
- § Some geometries re-optimized with MP2
- § Size of basis set: ~1,000 basis functions
- § Calculations performed at ERDC (T3E) and AHPCRC (T3E)



# TS For Polymerization of Ethylene





# RESULTS - NLO



Systematic investigation of excited state energies and oscillator strengths of free-base porphyrins, phthalocyanines, and their metal complexes, representing a broad range of  $\pi$ -conjugated systems, using time-dependent density functional theory (TD-DFT).






Figure 3.  $\pi$ -conjugated systems




# RESULTS - NLO



Computed  $T_1-T_n$  compared to experimental results


86 Experimental Values (in solution)  
Mean Absolute Error = 0.11  
Maximum Error = 0.31



Computed  $S_0-T_1$  compared to experimental results

47 Experimental Values (in solution)  
Mean Absolute Error = 0.14  
Maximum Error = 0.33

100,000 hours on SGI O2K, Compaq GS320, IBM SP/P3 at ASC



# SUMMARY

## High Energy Density Matter

- High-nitrogen/polynitrogen compounds are substantially more energetic than hydrazine.
- Trityldiazonium cation is not a stable polynitrogen precursor.
- Trimethylsilyldiazonium cation is marginally stable, replacement of methyls with fluorines increases stability.

## Polyhedral Oligomeric Silsesquioxanes (POSS)

- Nuclear quantum effects are important in proton transfer reactions -- lower barriers by about 4 kcal/mol for water-assisted condensation of trisilanol
- Most effective catalyst for olefin polymerization is  $T_4$  cage
- Barrier is still ~10 kcal/mol too high: Ti-POSS, Ti-siloxanes may not be good catalysts for this reaction

## NLO materials



- Time-dependent density functional theory accurately predicts NLA in free-base porphyrins, phthalocyanines, and their metal complexes.
- Mean absolute error of 0.11 eV for computed triplet-triplet excitation energies
- Mean absolute error of 0.14 eV for computed singlet-triplet excitation energies




# FUTURE DIRECTIONS

## HEDM

- Thermal decomposition mechanisms of energetic hydrocarbons (e.g., BCP).
- Energetic Ionic Liquids (QSPR models)





# ACKNOWLEDGEMENTS



**POSS:** Takako Kudo, Shawn Phillips, Simon Webb, Frank Feher, Joe Lichtenhan


**HEDM:** Jeff Mills, Greg Drake, Karl Christe, Bill Wilson, Ashwani Vij, Vandana Vij

**NLO:** Kiet Nguyen, Paul Day

**GAMESS:** Graham Fletcher

**MSRCs, DCs:** ASC, ARL, ERDC, NAVO, MHPCC, AHPCRC, ARSC, AFFTC


**CHSSI funding (CCM-2, CCM-4, MBD-01)**



# Backup Slides








# PROJECT OVERVIEW - NLO




## Technical issues being addressed using CCM

### 2. "Tuning" of absorption spectrum by benzannulation, halide substitution

System	Property						
	M	X	R	IP	S <sub>0</sub> -S <sub>n</sub>	S <sub>0</sub> -T <sub>1</sub>	T <sub>1</sub> -T <sub>n</sub>
PH <sub>2</sub>	H <sub>2</sub>	H	H	E, C	E, C	E, C	E, C
ZnP	Zn	H	H	E, C	E, C	E, C	E, C
TPPH <sub>2</sub>	Zn	H	φ	E, C	E, C	E, C	E, C
ZnTPP	Zn	H	φ	E, C	E, C	E, C	E, C
ZnTPPBr <sub>8</sub>	Zn	Br	φ	C	E, C	E, C	E, C



IP = Ionization Potential, S<sub>0</sub>-S<sub>n</sub> = Ground State Spectrum,  
 S<sub>0</sub>-T<sub>1</sub> = Singlet-Triplet Gap T<sub>1</sub>-T<sub>n</sub> = Triplet-Triplet Spectrum  
 E = Experiment, C = Calculated



# RESULTS - POSS


## Mechanism of formation: role of solvent ( $H_2O$ ) & substituents (R)


RSi(OH) <sub>3</sub> + RSi(OH) <sub>2</sub> Si(OH) <sub>2</sub> Si(OH) <sub>2</sub> + H <sub>2</sub> O			
R	Energy barrier (kcal/mol)		
	HF/6-31G*	MP2/6-31G†	
H	30.4 (16.7)	10.9 (-9.3)	
Me	28.2 (14.7)	7.7 (-13.3)	
<i>t</i> -Bu	34.3 (24.9)	9.8 (-9.3)	
Ph	31.1 (18.2)	7.9 (-16.4)	

Values in parentheses are for water-catalyzed results.

Values in parentheses are for water-catalyzed results.

Kudo, T., Gordon, M.S. J. Am. Chem. Soc., 120, 11432 (1998)  
Kudo, T., Gordon, M.S. J. Phys. Chem. A, 104, 4058 (2000)





# RESULTS - POSS




## Mechanism of formation

### Key steps


1. Hydrolysis of  $\text{RSiX}_3$  ( $\text{R}=\text{H}, \text{CH}_3, \text{t-butyl}, \text{etc.}; \text{X}=\text{Cl}$ )  
$$\text{RSiCl}_3 + \text{H}_2\text{O} \rightarrow \text{RSiCl}_2\text{OH} + \text{HCl}$$
$$\text{RSiCl}_2\text{OH} + \text{H}_2\text{O} \rightarrow \text{RSiCl(OH)}_2 + \text{HCl}$$
$$\text{RSiCl(OH)}_2 + \text{H}_2\text{O} \rightarrow \text{RSi(OH)}_3$$
2. Condensation of  $\text{RSi(OH)}_3$  to disiloxane  
$$2 \text{RSi(OH)}_3 \rightarrow \text{R(OH)}_2\text{SiOSi(OH)}_2\text{R} + \text{H}_2\text{O}$$
3. Condensation of disiloxane to  $\text{D}_3, \text{D}_4$   
$$\text{RSi(OH)}_3 + \text{R(OH)}_2\text{SiOSi(OH)}_2\text{R} \rightarrow \text{D}_3 + 2\text{H}_2\text{O}$$
  
$$[2+2]: 2\text{R(OH)}_2\text{SiOSi(OH)}_2\text{R} \rightarrow \text{D}_4 + 2\text{H}_2\text{O}$$
$$[3+1]: \text{RSi(OH)}_3 + \text{R(OH)}_2\text{SiOSi(OH)}_2\text{R} \rightarrow \text{D}_4 + 2\text{H}_2\text{O}$$

Ring Expansion:  $\text{RSi(OH)}_3 + \text{D}_3 \rightarrow \text{D}_4 + \text{H}_2\text{O}$
4. Condensation of  $\text{D}_3, \text{D}_4$  to POSS (in progress)  
$$2\text{D}_4 \rightarrow \text{T}_8 + 4\text{H}_2\text{O}$$


....




D3



D4




T8



# RESULTS - POSS




## Nuclear quantum effects in condensation reactions



Level	Monomer x 2	Reactive Complex	Transition State	Products	CPU Time
RHF/6-31G*	0.0	-10.4	30.4	-7.2	1.0
NEO-HF/2	0.0	-11.4	24.9	-5.9	1.1
NEO-HF/6	0.0	-12.6	24.7	-6.7	1.8

Units in kcal/mol



Hammes-Schiffer, S. J. Phys. Chem. A 102 (1998), 10443


Webb, S.P., Agarwal, P.K., and Hammes-Schiffer, S. J. Phys. Chem. B, 104(2000), 888

Webb, S.P. and Hammes-Schiffer, S. J. Chem. Phys. 113 (2000), 5214




# RESULTS - POSS


Molecular "sieves": preferential capture  $N_2$  vs.  $O_2$ ?



TS structure of  $T_8 + O_2$



$T_{10}$  and  $T_{12}$  calculations in progress  
HPC requirements: ~50,000 node-hrs, AHPARC T3E, 256 GB



# RESULTS - NLO




## B3LYP $S_0-T_1$ Excitation Energies (in eV)

System	6-31G(d)	Error	Exp
Porphyrin ( $1^3B_{2u}$ )	1.42	0.16	1.58 <sup>a</sup>
Zinc Porphyrin ( $1^3B_{1u}$ )	1.65	0.07	1.72 <sup>b</sup>
Tetraphenylporphyrin ( $1^3B_1$ )	1.31	0.14	1.45 <sup>c</sup>
Zinc Tetraphenylporphyrin( $1^3B_1$ )	1.53	0.06	1.59 <sup>d</sup>
Zinc Phthalocyanine ( $1^3B_{2u}$ )	1.05	0.08	1.13 <sup>e</sup>
Zinc Tetrabenzporphyrin ( $1^3B_{1u}$ )	1.41	0.16	1.57 <sup>f</sup>
Phthalocyanine ( $1^3B_{1u}$ )	1.18	0.06	1.24 <sup>g</sup>
Mean Error		0.10	


<sup>a</sup>Gouterman, Khalil, *J. Mol. Spectrosc.* 1974, 53, 88. (EPA (5:5:2) mixture of ethyl ether to isopentane to ethanol) and 50% ethyl iodide at 77 K) <sup>b</sup>Gradyushko, Tsvirko, *Opt. Spectrosc.* 1971, 31, 291.(EPA at 77 K) <sup>c</sup>Gouterman, Khalil, *J. Mol. Spectrosc.* 1974, 53, 88. (EPA at 77 K) <sup>d</sup>Walters et al., *J. Phys. Chem.* 1995, 99, 1166.(1:1 mixture of ether to ethanol at 77 K) <sup>e</sup>Vincett et al.,K. E. *J. Chem. Phys.* 1971, 55, 4131. (1-chloronaphthalene at 77 K) <sup>f</sup>Bajema, Gouterman, *J. Mol. Spectrosc.* 1971, 39, 421 (octane at 77 K) <sup>g</sup>McVie et al., *J. Chem. Soc. Faraday Trans. II* 1978, 74, 1870 (1-chloronaphthalene at 77 K)

Nguyen, K. A., Day, P. N., and Pachter, R., *J. Chem. Phys.*, 110 (1999) 9135  
Nguyen, K. A., Day, P. N., and Pachter, R., *J. Phys. Chem. A*, 103 (1999) 7378  
Nguyen, K. A., Day, P. N., and Pachter, R., *J. Phys. Chem.*, 104 (2000) 4755  
Nguyen, K. A. and Pachter, R., *J. Phys. Chem.*, 104 (2000) 4549



# RESULTS - NLO

## Comparison with Experiment: ZnPc



HPC Requirements: ~100,000 CPU hours, on SGI O2K + IBM SP3 + SGI O2K @ ASC